

Controlling wild oats

This trial is funded by the GRDC and is part of a collaborative project. It was conducted with Sam Kleemann, University of Adelaide and Peter Boutsalis, Plant Science Consulting.

Key findings

- One year of full wild oat control reduced the wild oat seedbank to 8 seeds per square metre in 2010.
- A selective post emergent herbicide or an early hay cut were the most effective strategies for reducing the wild oat seedbank.

Why do the trial?

The density of wild oats (*Avena fatua*) is increasing in the Mid North. This is due to an increase in cereal cropping intensity and the increase in herbicide resistance to Group A fop and dim herbicides. Also, traditional measures implemented for the control of annual ryegrass such as pre-emergent herbicides, export oaten hay, chaff carts and crop topping are generally less effective against wild oats.

This trial aims to evaluate the effect of long term management strategies on the wild oat seedbank and measure the efficacy of various control techniques. Specifically, the trial will demonstrate the value of single year and back-to-back years of seed set control, pre-emergent and post emergent herbicides, hay cutting and chaff cart for driving down the wild oat seed bank.

Herbicide resistance and wild oats – Peter Boutsalis, Plant Science Consulting

Herbicide resistance in wild oats occurs in all cereal growing regions. A random survey conducted in 1995 detected 5% of wild oat samples collected from NE Victoria as resistant to Hoegrass. In 2006, the number had increased to only 8% in a similar survey. In the Mid-North 35% of paddocks contain wild oat and of these 9% were resistant to Topik or Wildcat (Table 1).

Often wild oats can be resistant to certain Group A Fop herbicides and not others eg. resistant to Wildcat but not Verdict. In addition some fop-resistant wild oats are cross-resistant to Mataven, although Mataven may have never been used previously. Dim/Den herbicides can be effective on fop-resistant wild oats although this can be variable. About 50% of wild oats resistant to Topik or Wildcat are also resistant to Axial and / or Mataven.

A small number of Group B resistant wild oats have been reported. No resistance to IMI (Group B) chemistry or to trifluralin (Group D) or triallate (Group J) has been detected.

Table 1: Occurrence of herbicide resistance across South Australia and Victoria as detected by **random** sampling. Data is % of **paddocks** with herbicide resistant wild oats. Resistance is defined as samples where $\geq 20\%$ survival was detected in a pot test. A dash indicates no test with that herbicide.

Herbicide	Victoria Western (2005)	Victoria Northern (2006)	SA Mid North (2008)	SA Eyre Peninsula (2009)
Fields with wild oats	31%	81%	35%	36%
Hoegrass	17	8	>9	>2
Topik/Wildcat	-	-	9	2
Verdict	-	-	4	2
Axial/Achieve	-	2	6	2
Mataven	-	-	14	0
Atlantis	-	-	0	0

How was it done?

This trial was established in a grower paddock, north of Clare (White Hut) on an existing patch of wild oats in 2009. The majority of wild oat seed was within 2cm of soil depth, some being on the soil surface, and the oats are 100% susceptible to group A post emergent selective herbicides. The trial was established as a randomised complete block design with 3 replicates.

In 2009, the trial was sown to Catalina wheat and in 2010 Commander barley, and wild oat control treatments were applied to the same plots each year. The herbicides treatments were applied IBS (incorporated by sowing) prior to sowing with a commercial seeder (i.e. knife-point & press wheels).

Treatments:

- 1) nil
- 2) Trifluralin 1.5L/ha (incorporated by sowing - IBS)
- 3) Trifluralin 1.5L/ha and Avadex Xtra 2.0L/ha (IBS)
- 4) Trifluralin 1.5L/ha and Avadex Xtra 2.0L/ha (IBS) + Axial 200ml/ha (GS39)
- 5) Trifluralin 1.5L/ha (IBS) + early hay cut
- 6) Trifluralin 1.5L/ha (IBS) + chaff cart
- 7) Trifluralin 1.5L/ha (IBS – 2009 and 2010) + Axial 200ml/ha (GS39 2009 only)
- 8) Trifluralin 1.5L/ha (IBS – 2009 and 2010) + Axial 200ml/ha (GS39 2009 and 2010)

Wild oats were counted 4 to 6 weeks after sowing using a 20 cm x 30 cm quadrat from 4 random locations within each plot.

The initial seedbank at the site in 2009 was 400 wild oat seed per square metre to 10cm of soil depth and 150 plants per square metre emerged in the nil treatments after sowing.

The hay cut was performed at the beginning of the hay cutting season, and the chaff cart was simulated by removing wild oat heads at the beginning of harvest as determined by district practice in both cases.

Results

Clear differences in the wild oat seedbank have been shown for the different management strategies applied in 2009 and 2010 (Figure 1). With no control the wild oat seed density increased from 400 seeds per square metre in 2009 to 8092 seeds per square metre in 2011, a 20 fold increase. Similar increases in the wild oat seedbank were measured for trifluralin applied alone or when mixed with Avadex Xtra, which provided limited wild oat control.

When Axial was included as a late selective post emergent application the seedbank declined to less than 64% of the original 2009 level (400 seeds per square metre). This treatment may not be as effective on wild oats with resistance to group A herbicides.

One year of full wild oat control reduced the wild oat seedbank to 8 seeds per square metre in 2010. While the trial average was only 8 seeds per square metre, 19 wild oat plants per square metre was counted 4 weeks after sowing and without control meant the seedbank increased significantly in 2011. 2 years of full control has reduced the seedbank down to about 500 seeds per square metre, which is unexplainably higher than the initial seedbank of 400 seeds per square metre.

Of the cultural control practices the early hay cut was an effective strategy for reducing the wild oat seedbank. The cut done early and did not include raking or super conditioning, which might increase wild oat seed shed. The simulated chaff cart treatment was applied early in the harvesting window, but had limited success as many of the wild oats had already dropped seed by the time of harvest.

The success of these control treatments might also be influenced by the competitiveness of the crop, soil type, growing season rainfall and finish to the season. So, in seasons with a mild finish or later districts it is likely that more wild oat seed will be set.




Figure 1. The effect of different management strategies on pre-sowing (March) wild oat seed density at Clare in 2010 and 2011.

Acknowledgments

The Hart Field-Site Group wish to thank Andrew and Richard Hawker and Brian Jamieson for the use of their paddock and their cooperation with this trial work.