

Naparoo wheat grazing demonstration at Tungamah

Dale Grey¹, Sandy Quinlivan² and Alison Frischke³

¹ DPI Victoria, Cobram

² Formerly DPI Birchip

³ Birchip Cropping Group

Key point

- Grazing Naparoo wheat reduced grain yield by up to 19%.

Location: Tungamah, Victoria

Rainfall:

Annual: 550mm (avg 510mm)

GSR: 224mm (avg 334mm)

Soil:

Type: Brown loam over heavy grey clay

Sowing information:

Variety: Naparoo wheat

Sowing date: 17 April 2011

Sowing rate: 90kg/ha

Fertiliser: 110 kg/ha MAP; 12 kg/ha foliar nitrogen; 60 kg/ha nitrogen as urea

Treatments: Grazed (74DSE/ha) and ungrazed

Row spacing: 30cm

Paddock history:

2010 — canola

Plot size: 6m x 6m

Replicates: 3

Aim

To determine the effect of grazing on grain yield of Naparoo wheat.

Method

A dryland demonstration site was established at Tungamah to determine the impact of grazing on wheat yield. Naparoo wheat was sown into moisture on 17 April 2011 at 90kg/ha with 110kg/ha MAP on a 30cm row spacing in a 24ha paddock.

Naparoo was chosen for its high biomass production, rust resistance and its winter habit for grazing. The crop was top dressed on 18 May 2011 with 12kg/ha foliar nitrogen at GS14 (four leaf). Dry matter (DM) cuts were taken on 1 July 2011 at GS16.5 to determine available biomass.

The crop was grazed on 2 July 2011 by 400 ewes in lamb (38 DSE/ha) but removed on 3 July 2011 due to two ewe deaths. The ewes were lambd out in another paddock. The 400 ewes with 400 lambs (74 DSE/ha) were reintroduced on 25 July 2011 for three weeks. The crop was then top dressed on 15 August 2011 with 60kg/ha nitrogen as urea. The crop was grown to maturity and harvested for grain using a plot header.

Results

The ungrazed paddock contained 596kg/ha DM (see Table 1) at 6.5 leaf stage, 12 tillers, no nodes. Plant establishment was fair due to the dry autumn start with the crop looking thin but tillering well. The effect of grazing varied across the paddock. In two areas of the paddock the yield loss was marked, while in another area there was no yield difference between grazed and ungrazed plots. On average the yield loss was 19% (see Table 1) but this was not statistically significant across the trial. Protein levels were similar for grazed and ungrazed treatments.

TABLE 1 Dry matter and grain yield results for grazed and ungrazed Naparoo wheat

Treatment	Dry matter (kg/ha)	Yield (t/ha)	Screenings (%)	Protein (%)
Grazed	596	4.05	1.1	9.3
Ungrazed		4.98	0.7	9.2
		P = 0.193 lsd = 2.07 cv% = 13		



Photo: Dale Grey



Observations and comments

The sheep grazed an estimated 700kg of DM/ha during the three-week grazing period. The profitability of the paddock was not compromised by grazing and provided stock with feed during July.

Cereal crops grazed during the vegetative stage represent a high-quality feed source but are low in salt and high in potassium, which can reduce magnesium absorption in the rumen. Animals grazing cereal crops (especially wheat) require supplementation with a lick made of equal parts of magnesium oxide, ground limestone and salt. The limestone will provide calcium to pregnant and lactating ewes.

It is important to introduce animals to all new feeds gradually and provide supplementary hay. A gradual introduction will minimise the potential for scouring as the rumen bacteria adapt to the new feed source, while the hay will provide adequate fibre to assist rumen function. Careful introduction to feed on sunny days (avoid overcast conditions) and access to hay will also avoid nitrate poisoning when grazing oat and canola crops, particularly for more susceptible livestock such as heavily pregnant and lactating ewes and cows.

SPONSORS

This project is funded by the Grains Research and Development Corporation (GRDC) and Caring for our Country through the northern Victoria *Grain and Graze 2* (GRDC project BWB00018).

Thanks also go to farmer co-operators Josh and Jenny Buerckner.

CONTACT


Dale Grey

DPI Victoria, Cobram

T: (03) 5871 0600

E: dale.grey@dpi.vic.gov.au

Maximise your crop potential... Starfire™ from



"YOU'LL NOTICE A DIFFERENCE"

HUTCHEON & PEARCE

www.hutcheonandpearce.com.au

Albury 02 6058 6800 Condobolin 02 6895 2666 Cowra 02 6341 4800 Parkes 02 6862 1666 Temora 02 6977 1100 Wagga Wagga 02 6933 7900



JOHN DEERE

Precision Farming... Solutions to Eliminate vehicle drift, saving you time and money for years to come.

StarFire™ 3000 Receiver

The StarFire™ 3000 is a 66-channel, multi-frequency differential Global Navigation Satellite System (GNSS) receiver. The StarFire™ 3000 Receiver has access to additional satellite signals, true 3D integrated terrain compensation, and upgradeable accuracy. With the StarFire™ 3000 Receiver's enhancements, it provides more reliable positioning in areas with intermittent satellite availability or in other uncontrollable satellite environments. The StarFire™ 3000 also responds faster and more accurately to changes in terrain and has the ability to increase the level of accuracy without having to purchase a new receiver.

For more information contact your local John Deere dealer:

Hutcheon A&G Pty Ltd

ABN 26 096 546 682



Narrandera
(02) 6959 1522

Finley
(03) 5883 1655

Coleambally
(02) 6954 4280