


Trial number: NGA0702
Site: 'Oodnadatta' Moree
Co-operator: Hugh Ball

Planting date: 5/06/2007
Harvest date: 30/10/2007
PreDicta B crown rot result: 0 pg DNA/ g soil (Below detection limit)
In-crop rainfall: 103 mm

Impact on yield from addition of crown rot (CR)




★ = significant **reduction** in variety yield with addition of crown rot
 NSD = no significant difference in variety yield with addition of crown rot
 CV=16%, LSD (5%) = 172 kg/ha

With the addition of crown rot:

- Barley recorded an average 25% yield reduction (~280 kg/ha)
- Bread wheat recorded an average 35% yield reduction (~290 kg/ha)
- Bellaroi recorded a 77% yield reduction (~620 kg/ha)

Impact on screenings from addition of crown rot (CR)



★ = significant **increase** in variety screenings with addition of crown rot
 NSD = no significant difference in variety screenings with addition of crown rot

With the addition of crown rot:

- Barley recorded an average 4% **reduction** in screenings
- Bread wheat recorded an average 11% increase in screenings
- Bellaroi recorded a 21% increase in screenings

Key messages

Trial planted on very marginal soil moisture with extreme moisture stress pre and post flowering.

- Moderate to high crown rot yield loss situation
- Barley average yield ~1100 kg/ha with average bread wheat yield ~800 kg/ha
- Similar levels of absolute yield loss between barley and bread wheat
- EGA Wylie recorded the least impact from crown rot amongst the bread wheats
- No impact from crown rot on barley quality