

**Project:** Evaluation of fungicides for crown rot in wheat  
**Trial:** RD1114  
**District:** Condamine  
**Variety:** EGA Gregory  
**Sown:** 15/05/2011

	Application date	Crop stage
T1	22/07/2011	GS15
T2	5/08/2011	GS31-32

**Disclaimer:**

This document is based on the results from an individual trial and may contain experimental use patterns that are currently off-label. **This document does not provide any interpretation and should not be taken as an endorsement of any unregistered use pattern.**

Professional advice should be sought for specific recommendations to ensure access to the most up to date information and knowledge.

***Any product referred to in this document must be used strictly as directed, and in accordance with all label or permit instructions. Always consult the label prior to use.***

Trt No.	Product	Timing	Rate ml/ha	Harvest	Grain Quality			Pathology on Stubble		
				Yield kg/ha				Bipolaris %	CR Severity	Fusarium %
				3/11/2011	Test Weight kg/hL	Screenings %	Protein %			
1	Untreated	-	-	3221	77	3.4	9.0	12 b	3.8	7
2	Folicur	T1	145	3193	75	1.3	9.2	10 b	6.6	9
3	Folicur	T2	145	3366	75	1.4	10.0	23 a	5.5	15
4	Folicur	T1 & T2	145 x 2	3263	75	1.8	9.1	7 b	4.1	6
5	Folicur	T1	290	3347	76	3.3	8.9	8 b	5.9	9
6	Folicur	T2	290	3349	79	1.2	9.9	10 b	7.6	4
7	Gp 3 F	T1 & T2	290 x 2	3206	76	1.4	8.7	8 b	3.3	8
8	Prosaro	T1	300	3249	78	1.5	9.2	7 b	5.8	10
9	Prosaro	T2	300	3218	78	1.3	10.1	7 b	6.5	11
10	Prosaro	T1 & T2	300 x 2	3403	77	2.7	9.7	12 b	4.6	6
P =				0.81	0.87	0.36	0.48	0.09	0.87	0.89
LSD =				nsd	nsd	nsd	nsd	8.4 (10%)	nsd	nsd
CV =				6.2						