

Project: Botryosphaeria Mould in Wheat
Trial: RD1211
District: Inglestone
Crop: Wheat cv. EGA Gregory

Disclaimer:

This document is based on the results from an individual trial and may contain experimental patterns that are currently off-label. This document does not provide any interpretation and **should not be taken as an endorsement of any unregistered use pattern.**

Professional advice should be sought for specific recommendations to ensure access to the most up to date information and knowledge.

Any product referred to in this document must be used strictly as directed, and in accordance with label or permit instructions. Always consult the label prior to use.

Timing	Date	Wheat growth stage
T1	10/09/2012	GS 58 - 66 (Ave = 63)
T2	21/09/2012	GS 71 - 83

Trt No.	Product	Rate ml/ha	Timing	Adjuvant	Yield 17/10/2012 kg/ha
1	Untreated	-	-	-	2852
2	Tilt	500	T1 (start of flowering)	-	2836
3	Tilt Xtra	500	T1	-	2739
4	Folicur	290	T1	-	2948
5	Amistar Xtra	400	T1	Adigor 2%	2792
6	Opera	500	T1	-	2647
7	Prosaro	300	T1	-	2861
8	Prosaro	300 x 2	T1 + T2	-	2680
9	Tilt	500	T2 (11 days after T1)	-	2682
10	Tilt Xtra	500	T2	-	2655
11	Folicur	290	T2	-	2844
12	Amistar Xtra	400	T2	Adigor 2%	2687
13	Opera	500	T2	-	2896
14	Prosaro	300	T2	-	2657
P =					0.77
LSD =					nsd
CV =					8.56

Field comments:

Trial conducted on a property that had experienced Botryosphaeria in previous 2 seasons

No rain during flowering period

No disease detected

No yield differences