

West Midlands Group 2015 Research Annual Template

Ripping by time of nitrogen demonstration

Bill Bowden and Chris Wilkins, WMG

Purpose: To demonstrate that, in the right season, ripping affects nitrogen uptake efficiency and availability to crops

Location: Charles Roberts, "Kayanaba", Dandaragan

Soil Type: Wakea, red loamy sand

Soil Test Results:

Soil analyses for adjacent Summit K trial site												
Depth	NH4 +	NO3-	P	K	S	Cu	Zn	Org C	pH[Ca]	Al	EC	PBI
0-10	14	2	29	38	6	0.64	0.3	0.69	5.1	0.6	0.04	35
10-20	8	2	19	32	5	0.63	0.15	0.59	4.6	0.3	0.02	44
20-30	3	1	8	32	6	0.31	0.02	0.31	4.5	1.2	0.02	45

Rotation: Canola stubble and burnt windrows. 2014: canola, 2013: wheat, 2012: oats

Growing Season Rainfall (April- October 2015): as for 2015 the WMG field day site

BACKGROUND SUMMARY

Removing chemical (acidity) or mechanical (plough or traffic) pans allows crop roots to penetrate soils more rapidly and so keep up with leaching nitrate nitrogen and improve nitrogen uptake efficiency. Ripping or deep cultivation can also stimulate nitrogen mineralization. Deeper roots also allow crops to access deeper water and so perform better in seasons with a dry finish or terminal drought. Given the right shaped season this work will demonstrate some of those effects.

We also tested direct diagnostics at this site using a long ripped strip and canola windrow effects

TRIAL DESIGN

This is a split, split plot latin square, design. Ripped and non-ripped plots are crossed with 4 time of nitrogen treatments (nil, 50 kg N/ha at 4 weeks after seeding, 8 weeks after seeding and at 4 and 8 weeks after seeding..

Plot size: 10 metres by 3 metres

Machinery use: Ripped with the DAFWA Merredin ripper to 35 cm with modified box boots. The trial was sown by the farmer as a bulk wheat crop across the ripped treatments

Repetitions: 4

Crop type and varieties used: Mace wheat


Seeding rates and dates: Sown on 25 May with Mace wheat at 100kg/ha. 80 kg/ha of MAXamRITE and 50 litres/ha of MAXamFLO applied at seeding. 100kg NKS applied in late June. Rates of basal nutrients- 50kg N/ha, 15 kg P/ha, 22 kg K/ha and 9 kg S/ha

Treatment rates and dates: (The 4/5 WAS N treatments were applied at 110 kg urea/ha on 1 July and the 8/9 WAS N treatments were applied at 110 kg urea/ha on 29 July).

Herbicide rates and dates:

First knock (pre seeding)	Seeding 25th May	Post seeding	Post seeding 300mL
80L Water Rate	80L Water rate	100L Water rate	Tebuconazole
0.4L 2,4-D ester 680	2L trifluralin 480	0.8L 2,4-D ester 680	Alpha Cyper 125mL
1.8L Roundup 570	1L Sprayseed	10g Logran	Wetter 1000 0.2%
	300ml chlorpyrifos	50g Lontrel	

Other applications/ treatment rates and dates:



RESULTS/STATISTICS

The results from measurements during the season and at maturity are shown below.

		plt/M ²	Biomass in t/ha				Yield components				
		10 July	29 July	11 Aug	8 Sept	9 Nov	Yield	HI	head #	tgw	screen%
Mean	Rip	150	0.7	1.2	3.3	4.9	2.0	0.41	248	31.0	17
Mean	No rip	85	0.4	0.8	2.4	3.3	1.3	0.39	194	31.0	23
C of V	Rip	NA	0.25	0.2	0.10	0.11	0.09	0.05	0.11	0.05	0.25
C of V	No rip	NA	0.36	0.4	0.28	0.35	0.39	0.17	0.22	0.05	0.43

Results were averaged across N treatments for all in-season measurements. The final biomass and yield component measurements were made on the control (no N except for farmer basal applications) and the 100 kg N/ha split application treatments only.

The diagnostic ripped strip was sampled in two places in the paddock– (1) in a severely non-wetting area south of the trial where there was a marked response in both establishment and growth and (2) north of the trial where there were no establishment differences but a small growth response. The diagnostic windrows were sampled north of trial (2) on 10 July and also south of trial (3) on better crop performing country in the valley on heavier soil on 18th August (3). These were paired on/off samples with no replicates – see the table below.

Please complete and return to Anne Wilkins, West Midlands Group by **Thursday 4 February 2016**
 E: anne@wmgroup.org.au T: 08 9651 4008 F: 08 9651 4107 M: 0438 924 208

Rip strip yield components - Kanyana 2015										
		plants	spring tops	mature tops	head #	grain yield	HI	screen	tgw	grains
		#/M^2	kg/ha	kg/ha	/ M^2	kg/ha	total	<2mm %	grams total	#/hd total
sandy soil										
2 wettable	rip	166	3600	6700	305	2775	0.41	9.9	29.9	30
2 wettable	norip	120	3100	5700	285	2250	0.39	12.2	26.6	30
1 non wet	rip	105	2700	4700	210	2000	0.43	11.3	35.1	27
1 non wet	norip	66	1600	2200	168	775	0.35	61.3	23.6	20

Windrow yield components - Kanyana 2015										
		K %	tops dwt kg/ha		head #	yield	HI	<2mm	tgm	#/hd
		early	early	mature	/ M^2	kg/ha	total	%	gms	total
soil										
2 sandy	on	4.0	519	2088	205	1950	0.47	7.7	30.3	31
2 sandy	off	2.2	139	2150	250	1625	0.38	26.2	28.9	23
3 loamy	on	2.8	1744	10930	393	4600	0.42	1.6	31.8	37
3 loamy	off	1.4	1500	9700	470	4255	0.44	3.6	25.6	35

OBSERVATION/ DISCUSSION/ MEASUREMENTS

There has been a major response to ripping but little response to nitrogen above and beyond the farmer dressings. The ripping treatment markedly improved crop emergence numbers and timing. The very dry nature of the season after seeding has meant no leaching of nitrogen and therefore no improvement in nitrogen uptake efficiency due to faster root penetration on the ripped plots. Plant counts, estimated biomass levels and yield components are shown in the table of results, above.

There was a marked establishment and growth response to ripping and variability was much more marked on the non-ripped plots than on those which were ripped. Measures of surface soil wettability showed no difference between ripping so the better establishment was probably due to better and more uniform wetting on the rougher, more disturbed, ripped soil. Ripping gave better early growth but also better finishing conditions as reflected in higher harvest indices and lower screenings. The ripped strip samplings emphasized this point so it seems that the ripping allowed better access to stored sub-soil moisture over the harsh finish to the season. Plant samples from on and off the ripped strips showed no difference in K status.

We sampled on and off visually obvious windrows north of trial (2) on 10 July and south of trial (3) on heavier country on 18th August. Both sites showed unambiguous K deficiency which again was reflected in better grain size and lower screening measures on the K adequate windrow. The sandy site had more severe K deficiency and bigger responses than the loamy site which had a higher soil K status. Both areas would respond to K fertilizer applications.

PEER REVIEW/REVIEW

ACKNOWLEDGEMENTS/ THANKS

Thanks to GRDC for funding the project, Charles Roberts for use of the land and to Caz Abbey for technical assistance and dangerous videotaping.