



Sowing dates – getting the best from our varieties and optimising whole-farm wheat yield

2012 Trial Site



Project Partners



Jointly Funded by the

GRDC National Water-Use Efficiency Initiative and
DAFF National Adaptation and Mitigation initiative

GRDC
Grains
Research &
Development
Corporation
Your GRDC working with you


Australian Government
Department of Agriculture,
Fisheries and Forestry

Sowing dates – getting the best from our varieties and optimising whole-farm wheat yield

James Hunt (CSIRO), Neil Fettell (University of New England), Jon Midwood (SFS), Annieka Paridaen (SFS), Julianne Lilley (CSIRO), Paul Breust (FarmLink Research), Rohan Brill (NSW DPI), Brad Rheinheimer (CSIRO), John Kirkegaard (CSIRO)

Key Messages

- Very slow or slow maturing wheat varieties sown mid-late April and managed for high HI on average yield 0.8 t/ha more than mid-fast varieties sown mid-May.
- Simulations show that including a slow maturing variety in a farm program greatly increases average farm wheat yield (13-47%) and reduces risk.
- Know the optimal flowering window in your location and how to achieve it with different varieties from different sowing dates to optimise whole-farm wheat yield.
- Yield penalties for sowing too late are generally higher than for sowing too early, even at frosty sites.

Key Words

time of sowing, wheat, water-use efficiency

Background

In southern Australia, April-May rainfall has declined due to movement in the position of the sub-tropical ridge, a consequence of ozone depletion and/or global warming due to greenhouse gas emissions (<http://www.grdc.com.au/Media-Centre/Ground-Cover/Ground-Cover-Issue-101/Expanding-subtropic-zone-the-clue-to-dry-autumns>). Rain during this time is critical for establishing the currently popular mid-fast spring wheat varieties so that they flower at a time optimal for yield. Also, larger farm sizes and cropped area mean that contemporary sowing programs often exceed the available sowing opportunities. Extreme weather during spring has made achieving timely flowering of cereal crops increasingly critical to yield and farm profitability.

An opportunity exists to overcome this challenge by using rain which falls during the summer and early autumn fallow period to establish wheat crops much earlier than is currently practiced. At some locations in the region, the proportion of rain falling during the summer fallow period has increased, and growers are placing more emphasis on conserving the resource through complete control of summer fallow weeds and wide adoption of no-till seeding


systems. No-till allows much earlier planting into more marginal moisture and in response to smaller rainfall events, and deeper ('moisture seeking'). However, planting currently grown varieties in early autumn incurs unacceptable frost risk, and for this change to happen slow maturing varieties with either stronger vernalisation (winter wheats) or photo-period requirements (slow maturing spring wheats) will be necessary. Fortunately such varieties exist, but are currently undervalued and rarely used by producers as they are sown too late in national variety trials for their yield and quality potential to be expressed.

The aim of this set of experiments was to see how slow maturing wheats sown early into stored soil water yielded in comparison to mid-fast varieties sown during the usual sowing window.

Method

APSIM simulations 1890-2009 were used with a multiplier on yield for frost and heat damage to identify optimal flowering windows at four different locations with contrasting climates (Lake Bolac Vic – high rainfall, Temora/Junee NSW – medium rainfall and Condobolin NSW – low rainfall). Optimal flowering dates in each environment were 23 October at Lake Bolac, 28 September at Temora and 16 September at Condobolin (Figure 1).

Figure 1. Mean flowering date (Z65) and yield at each location from 120 year (1890-2009) APSIM simulation with a multiplier applied to yield for heat and frost.



In each location, experiments were established with commercially available and locally adapted spring milling wheat varieties of varying maturity sown at different dates such that they flowered on the same optimal date identified by APSIM. Maturity groups were classified as 'very slow' (Forrest at Lake Bolac and EGA Eaglehawk at Temora/Junee and Condobolin), 'slow' (Bolac at all sites), 'mid' (Derrimut at Lake Bolac and EGA Gregory at Temora/Junee and Condobolin), 'fast' (Lincoln at all sites) and 'very fast' (Axe at Condobolin 2011 only). Sowing date of the different maturity classes required to achieve optimal flowering date was consistent across environments. Optimal sowing date for the very slow maturity group was 15 April

with optimal sowing dates for subsequent groups being progressively ten days afterward (slow – 25 April, mid – 5 May, fast – 15 May, very fast 25 May). Experimental sowing dates deviated from these somewhat, but flowering of the different maturity classes largely coincided at all locations (Table 1).

We also applied regionally relevant management treatments intended to avoid excessive vegetative growth and increase HI, which included very low plant densities (Temora/Junee, Condobolin, Lake Bolac), plant growth regulators (PGRs - Lake Bolac and Temora 2011), defoliation (to simulate grazing) at Z30 (Junee and Condobolin 2012) and sub-soil manuring (Lake Bolac 2011).

Table 1. Experimental sowing and anthesis dates (Z65 – 50% of ears flowered) for the different maturity groups at each location in 2011 and 2012.

Maturity Group	Lake Bolac		Temora		Condobolin	
	Sowing Date	Anthesis Date	Sowing Date	Anthesis Date	Sowing Date	Anthesis Date
2011						
Very slow	15 April	19 October	15 April	2 October	15 April	15 September
Slow	27 April	16 October	27 April	3 October	27 April	13 September
Mid	6 May	14 October	9 May	5 October	5 May	13 September
Fast	20 May	20 October	19 May	4 October	16 May	16 September
Very fast	-	-	-	-	25 May	20 September
2012						
Very slow	18 April	24 October	18 April	14 October	19 April	28 September
Slow	27 April	13 October	26 April	11 October	30 April	24 September
Mid	8 May	13 October	8 May	13 October	9 May	28 September
Fast	18 May	18 October	17 May	13 October	18 May	24 September

Results

At Temora in 2011 there was a large yield advantage (0.8 t/ha) from sowing a very slow variety early (EGA Eaglehawk 15 April) and reducing seeding rate to improve HI (6.3 t/ha) compared to sowing mid and mid-fast varieties in their optimal window (EGA Gregory 9 May 5.4 t/ha, Lincoln 19 May 5.5 t/ha – Table 2) as currently practiced. There was a yield reduction in mid and mid-fast varieties sown at low densities and there was no effect of PGRs on yield. The yield benefit of the slow varieties may be an under-estimate of the value of early sowing

at this site. Seed bed moisture was perfect on 15 April following 16 mm of rain on 10 April, but was becoming marginal on 27 April and by 9 and 19 May irrigation (8 mm applied into press-wheel furrows with drippers) was required to establish the crop. Therefore in a dry-land farm situation where irrigation would not be possible, the early-sown slow varieties would have had an even greater yield advantage over the later-sown faster varieties as the rain that allowed them to emerge did not fall until 25 May.

Table 2. Grain yield and harvest index of four wheat varieties of different maturity sown at two plant densities at Temora in 2011 to flower on the same date.

Variety and sowing date	Grain yield (t/ha)		Harvest index (%)	
	40 plants/m ²	100 plants/m ²	40 plants/m ²	100 plants/m ²
EGA Eaglehawk (15 April)	6.3	6.0	0.41	0.39
Bolac (27 April)	5.9	5.7	0.42	0.39
EGA Gregory (9 May)	5.0	5.4	0.44	0.43
Lincoln (19 May)	4.8	5.5	0.44	0.44
P-value	0.009		0.018	
LSD (p=0.05)	0.5		0.01	

Results cont.

At Junee in 2012 there was again a large yield advantage (2.1 t/ha) from sowing a very slow variety early (EGA Eaglehawk 18 April 6.1 t/ha) compared to sowing mid and mid-fast varieties in their optimal window (EGA Gregory 8 May 5.2 t/ha, Lincoln 17 May 4.0 t/ha – Table 3). Defoliation generally reduced yields, with the exception of Bolac at 100 plants/m² and Lincoln at both densities

where the difference was not significant. Reduced plant density generally did not affect yield, but provided a small increase for un-defoliated Bolac and Lincoln. Time of sowing data show that there is generally less yield penalty from sowing earlier than is optimal, despite both sites experienced frosts around anthesis (Figures 2 and 3). At Temora in 2011, the highest yields achieved at each time of sowing had 5% frost damage (Figure 2).

Table 3. Grain yield of four wheat varieties (un-defoliated or defoliated) of different maturity sown at Junee in 2012 to flower on the same date. .

Variety and sowing date	50 plants/m ²		100 plants/m ²	
	Not defoliated	Defoliated	Not defoliated	Defoliated
EGA Eaglehawk (18 April)	5.9	5.6	6.1	5.4
Bolac (26 April)	5.8	5.4	5.5	5.3
EGA Gregory (8 May)	5.1	4.7	5.2	4.6
Lincoln (17 May)	4.3	4.1	4.0	4.1
P-value	0.034			
LSD (p=0.05)	0.3			

In 2011 at Condobolin, dry conditions in late winter reduced grain set in the earliest sowings and similar yields were obtained from sowing dates ranging from 15 April until 16 May. The higher biomass from sowing earlier was offset by a lower HI (Table 4), even

at a low plant density. Whilst there was an increase in HI and yield at low plant densities for slow maturing varieties sown early (EGA Eaglehawk, Bolac), there was a yield penalty at low plant densities for fast maturing varieties sown later (Lincoln, Axe).

Table 4. Grain yield and harvest index of five wheat varieties of different maturity sown at two plant densities at Condobolin in 2011 to flower on the same date.

Variety and sowing date	Grain yield (t/ha)		Harvest index (%)	
	30 plants/m ²	90 plants/m ²	30 plants/m ²	90 plants/m ²
EGA Eaglehawk (15 April)	3.4	3.1	0.37	0.32
Bolac (27 April)	3.3	2.9	0.38	0.35
EGA Gregory (5 May)	3.6	3.2	0.44	0.39
Lincoln (16 May)	2.8	3.0	0.46	0.45
Axe (25 May)	2.1	2.6	0.45	0.44
P-value	0.029		0.027	
LSD (p=0.05)	0.4		0.02	

At Condobolin in 2012, the highest yield using the traditional seeding rate was achieved from sowing Bolac on 30 April. Biomass production from the 19 April sowing was equally high but was negated by a much lower HI (0.26 cf. 0.32). This could be largely offset by a lower seeding rate under which biomass was not reduced but HI was raised to 0.29. Defoliation at Z30 was also successful in raising HI values at all sowing dates but this was countered by lower maturity biomass at all but the latest sowing date. At Lake Bolac in 2011, annual ryegrass resistant to herbicide groups A & B overran the first time of sowing and all low plant density treatments and these have been excluded from analysis. The slow variety Bolac sown on 27 April yielded 7.0 t/ha, the mid variety Derrimut sown on 6 May yielded 6.6 t/ha and the fast variety Lincoln sown on 20 May


yielded 6.0 t/ha (P=0.006, LSD(P=0.05)=0.6 t/ha).

At Westmere in 2012, the highest yields were achieved by the slow variety Bolac sown at its optimal time on 27 April and also at 8 May (Figure 2). The very slow variety Forrest flowered later than expected at its optimal time of sowing (Table 1), and did not yield well. However, seed bed moisture at 18 April was poor, and establishment was only 58 plants/m² at that time of sowing compared to 122 plants/m² at the 18 May sowing (target density was 160 plants/m²). Bolac and Derrimut sown 1 May were also the highest yielding treatments in a scaled down version of the experiment at Inverleigh (Figure 3).

Why do early sown crops yield more?

Early sown crops yield more in situations where the soil profile fills with water because roots grow deeper and use more water (Figure 2). Wheat roots grow at approximately 12 mm/day from germination to anthesis, and early sowing of slow maturing wheats increases this period and thus potential rooting depth. varieties as the rain that allowed them to emerge did not fall until 25 May. classes largely coincided at all locations (Table 1).

Figure 2. Grain yield of different varieties at Temora in 2011 sown at 100 plants/m² with no PGRs. Numbers next to data points are % frost-induced sterility.



Early sown crops also develop their canopy faster (Figure 3) and as a result lose less water to evaporation, and thus water is converted to dry-matter more efficiently (Table 5). Also the longer stem elongation phase of early sown slow maturing varieties mean that they intercept more radiation and grow more during this period, which increases grain number and thus yield (Figures 4 and 5).

Figure 3. Grain yield of different varieties at Junee in 2012 sown at 100 plants/m² and not defoliated. Frost induced sterility data not available at time of publication.




Table 5. Crop water-use (evaporation + transpiration), dry matter at maturity (Z92), WUE for dry matter and estimates of transpiration and evaporation for varieties of differing maturity sown in their optimal window at June in 2012.

Variety and sowing date	Water-use (mm)	Total dry matter (kg/ha)	WUE for dry matter (kg/ha.mm)	Transpiration* (mm)	Evaporation (mm)
Eaglehawk 18 April	345	11756	34	214	132
Bolac 26 April	348	12556	36	228	120
Gregory 8 May	344	9798	28	178	166
Lincoln 17 May	321	9229	29	168	153

*Transpiration estimated assuming a transpiration efficiency for dry matter of 55 kg/ha.mm




Figure 4. The relationship between growth during stem elongation and grain yield at Lake Bolac (●), Temora (○) and Condobolin (□) in 2011.




Figure 5. Phase durations for EGA Eaglehawk sown 15 April and Lincoln sown 19 May at Temora in 2011.

Advantages at the whole farm

Including a slow maturing wheat variety in a farm program increases the frequency of planting opportunities, and allows more crop to be sown and flower on time, increasing average farm yield. We conducted APSIM simulations over 50 years to compare sowing strategies at hypothetical farms at 5 sites in south-eastern Australia. On each hypothetical farm we assumed that the total area of wheat sown in each year takes twenty days to sow (e.g. 20 paddocks each taking 1 day to sow). Other crops were not considered in the simulation, and nitrogen, weeds, disease etc. were assumed not to limit yields, but a reduction for frost and heat damage based on screen temperatures was applied. Yields presented are farm average wheat yield across 50 years, for 20 paddocks sown according to the following strategies.

1. **Mid-fast varieties only.** In this scenario only varieties from the mid-fast maturity group (e.g. Scout, Mace, Lincoln, Spitfire) were sown. Sowing started on 5 May at the earliest and only if there was seed-bed moisture (i.e. after the autumn break). If the break had not arrived by 25 May, sowing started on that date and finished on 13 June regardless of

seed bed moisture.

2. **Slow + mid-fast varieties.** In this scenario two varieties of differing maturity were grown on the farm – very slow (e.g. Wedgetail, Forrest, Eaglehawk – sown 10 April to 4 May), and mid-fast (e.g. Scout etc. – 5 May onward). Each variety was sown within its appropriate window, but only if there was seed-bed moisture, and in the case of the slow variety, only if there was also 25 mm of stored soil water in the top 25 cm of the profile. If the break had not arrived before 25 May, sowing started on that date and finished on 13 June and only the mid-fast variety was sown. In this scenario the area sown to each variety changed from year to year based on rainfall pattern e.g. if the break fell by 15 April, 100% of the farm was sown to the slow variety. If the break fell on 5 May or later, only the mid-fast variety was sown.

At all locations average farm yield was significantly increased and production risk reduced if slow varieties were included in the sowing program (Table 6).

Table 6. Including a very slow maturing wheat variety in a farm program that allows early sowing increases average farm yield and reduces risk. Results are from APSIM simulation 1962-2011 with a frost and heat multiplier for yield and assume a farm wheat program takes 20 days to sow.

Strategy	1. Mid-fast varieties only – sowing window opens 5 May		2. Slow + mid-fast variety – sowing window opens 10 April				
	Average farm wheat yield (t/ha)	Paddock yields <1.0 t/ha (%)	Average farm wheat yield (t/ha)	Paddock yields <1.0 t/ha (%)	Average farm wheat yield increase (%)	Years in which slow variety sown (%)	Average area of slow variety sown (%)
Location							
Condobolin, NSW	1.5	48%	2.2	34%	42%	71%	59%
Temora, NSW	2.9	22%	3.8	14%	29%	78%	66%
Kerang, Vic	2.1	35%	3.0	22%	44%	49%	37%
Lake Bolac, Vic	4.5	4%	5.0	5%	13%	98%	84%
Morchard, SA	1.6	44%	2.3	32%	47%	29%	19%

Whilst growers may cringe at the prospect of keeping seed of multiple varieties on hand to sow an unknown area of crop each year, the yield benefits are substantial. The % area which was sown to the slow variety was often high, indicating that sowing opportunities for slow maturing wheats are quite frequent. The analysis also highlights a significant opportunity for breeding companies willing to focus on development of slow maturing wheats of milling quality, particularly photoperiod insensitive winter wheats.

by early sowing e.g. barley yellow dwarf virus (BYDV), wheat streak mosaic virus, Septoria tritici, take-all. Seed-dressed imidicloprid (Gaucho, Emerge) protects against BYDV, and flutriafol in-furrow (Impact, Intake) protects against take-all and Septoria tritici. Forrest has putative resistance to WSMV.

Putting it into practice

If planning to sow early, the most important thing is to know the optimal flowering window in your location and how to hit it using different varieties from a broad range of sowing dates. There are three basic strategies for achieving an optimal flowering time;

1. Plant winter wheats from early-March to mid-April
2. Plant slow maturing spring wheats from early to late April
3. Plant mid-fast varieties from late April into May

The **first two strategies require some stored soil water** to be present (25 – 30 mm plant available water – use a shovel to check soil is wet to at least 30 cm on most soil types). **These crops must germinate and emerge when they are sown.** Winter and slow maturing spring wheats will flower too late if they do not establish before the end of April. Choose paddocks that are relatively weed-free and do not require a good knock-down for grass weed control. If sowing early into high soil N, reduce seeding rates.

Best-bet milling wheat varieties for early sowing in southern NSW are listed in Table 8.

Be aware that some diseases can be exacerbated

Table 8. Best-bet milling wheat varieties for early sowing in southern NSW.

Variety	Sowing window in southern NSW	Yield in early sown NVT southern NSW (t/ha)		Quality in southern NSW	Disease Vulnerabilities	Comments
		West	East			
Bolac	15 April – 1 May	3.2	3.5	APH	Septoria tritici, leaf rust, yellow leaf spot	The highest yielding early-sown hard variety in southern NSW. Excellent stripe rust resistance, but watch leaf rust and YLS in a warm, wet finish. Small grain size and free tillering make it very cheap to sow but screenings can be an issue if sown too late.
EGA_Bounty	20 April – 5 May	3.0	3.3	AH	Yellow leaf spot, Septoria tritici (?)	Faster than Bolac but slower than Gregory in southern NSW so useful for late April/early May. More susceptible to lodging than all other varieties listed here other than Sunvale.
EGA_Eaglehawk	10 April – 25 April	3.0	3.3	APH	Stripe rust, Septoria tritici	Very slow maturing photoperiod sensitive spring wheat that will flower at the same time as Wedgetail from a mid-April sowing but hit Z30 ~3 weeks earlier, therefore not as suited to grazing.
EGA_Wedgetail	15 March – 1 May	2.5	3.1	APH	Stripe rust, yellow leaf spot	The dual purpose standard and an excellent grain-only option. Vernalisation requirement gives it a very wide sowing window, but beware wheat-streak mosaic virus (WSMV) on the slopes if sowing before mid-April. NVT results prior to 2011 when fungicides began to be used will underestimate yield potential due to stripe rust.
Forrest	15 April – 1 May	NA	3.5	ASW provisional	Septoria tritici	Photoperiod sensitive spring wheat, faster than Eaglehawk but slower than Bolac. It has useful resistance to WSMV and all three rusts. Better grain size than Bolac. It will have potential in southern NSW provided it achieves AH classification.
Sunvale	15 April – 1 May	3.0	3.3	APH	Yellow leaf spot, Septoria tritici	Similar maturity to Bolac in southern NSW and whilst it does not yield as well it is less prone to screenings and has better sprouting tolerance. It is free tillering so can be sown light, and it attracts no EPR so it is a good low-input option.
Wylah	15 March – 1 May	2.7	2.9	AH	Leaf rust, stripe rust, yellow leaf spot	Photoperiod insensitive winter wheat and therefore faster to flower than Wedgetail making it more suited to lower rainfall environments.